

Contents

Volume 101 (4) 1992

Melan, M. A. and Sluder, G. Redistribution and differential extraction of soluble proteins in permeabilized cultured cells: implications for immunofluorescence microscopy	731-743	Chevrier, V., Komesli, S., Schmit, A.-C., Vantard, M., Lambert, A.-M. and Job, D. A monoclonal antibody, raised against mammalian centrosomes and screened by recognition of plant microtubule organizing centers, identifies a pericentriolar component in different cell types	823-835
Takeda, T., Yamashita, Y., Shimazaki, S. and Mitsui, Y. Histamine decreases the permeability of an endothelial cell monolayer by stimulating cyclic AMP production through the H ₂ -receptor	745-750	Stephens, R. E. Tubulin in sea urchin embryonic cilia: post-translational modifications during regeneration	837-845
Leitch, A. R., Mosgöller, W., Shi, M. and Heslop-Harrison, J. S. Different patterns of rDNA organization at interphase in nuclei of wheat and rye	751-757	King, K. L., Stewart, M., Roberts, T. M. and Seavy, M. Structure and macromolecular assembly of two isoforms of major sperm protein (MSP) from the amoeboid sperm of the nematode, <i>Ascaris suum</i>	847-857
Solovei, I., Gaginskaya, E., Allen, T. and Macgregor, H. A novel structure associated with a lampbrush chromosome in the chicken, <i>Gallus domesticus</i>	759-772	Ramachandrala, A., Tiku, K. and Tiku, M. L. Tripeptide RGD-dependent adhesion of articular chondrocytes to synovial fibroblasts	859-871
Stuurman, N., de Graaf, A., Floore, A., Josso, A., Humbel, B., de Jong, L. and van Driel, R. A monoclonal antibody recognizing nuclear matrix-associated nuclear bodies	773-784	Lu, M. L., McCarron, R. J. and Jacobson, B. S. Initiation of HeLa cell adhesion to collagen is dependent upon collagen receptor upregulation, segregation to the basal plasma membrane, clustering and binding to the cytoskeleton	873-883
Liu, D. F., El-Alfy, M. and Leblond, C. P. Visualization of DNA within mitochondria by osmium-ammine staining of mouse duodenal crypt cells	785-793	Szekanecz, Z., Humphries, M. J. and Ager, A. Lymphocyte adhesion to high endothelium is mediated by two β_1 integrin receptors for fibronectin, $\alpha_4\beta_1$ and $\alpha_5\beta_1$	885-894
Saulnier-Michel, C., Fromont-Racine, M. and Pictet, R. Co-expression of insulin and somatostatin genes in pancreatic endocrine cells selected for their high level of insulin gene transcription	795-799	Schenkman, S. and Mortara, R. A. HeLa cells extend and internalize pseudopodia during active invasion by <i>Trypanosoma cruzi</i> trypomastigotes	895-905
Marks, J., Fankhauser, C. and Simanis, V. Genetic interactions in the control of septation in <i>Schizosaccharomyces pombe</i>	801-808	Cannon, G. J. and Swanson, J. A. The macrophage capacity for phagocytosis	907-913
Steffen, W. and Linck, R. W. Evidence for a non-tubulin spindle matrix and for spindle components immunologically related to tektin filaments	809-822	Index of Authors and Titles	915-918
		Subject Index	919-926